arXiv:1302.3052v1 [gr-qc] 13 Feb 2013

# Marginally outer trapped surfaces in higher dimensions

Tim-Torben Paetz and Walter Simon

Gravitationsphysik, Fakultät für Physik, Universität Wien
Boltzmanngasse 5, A-1090 Wien, Austria

**Abstract**

We review the basic setup of Kaluza-Klein theory, namely a 5-dimensional vacuum with a cyclic isometry, which corresponds to Einstein-Maxwell-dilaton theory in spacetime. We first recall the behaviour of Killing horizons and its generators under bundle lift and projection. We then show that the property of compact surfaces of being (stably) marginally trapped is preserved under lift and projection provided the appropriate (”Pauli-”) conformal scaling is used for the spacetime metric. We also discuss and compare recently proven area inequalities for stable axially symmetric 2-dimensional and 3-dimensional marginally outer trapped surfaces.

## 1 Introduction

The basic setup of Kaluza-Klein (KK) theory consists of a 4+n dimensional manifold with n Killing vectors, which can be regarded as a principal fibre bundle over a 4d base (see e.g. [1, 2]). In 5 dimensions in particular, stationary black holes have been studied extensively and revealed much richer structures than in 4d (cf e.g. [3]–[8]). There is, however, a more subtle motivation for studying black holes in (4+n) dimensions: Suitable adaptions of Hawking’s rigidity theorem assert the existence of n+1 spacelike isometries on generic (non-static) stationary event horizons [9], and under suitable technical assumptions, such Killing fields can be extended off the horizon. From this point of view, the higher dimensional black hole is the ”fundamental” object which by itself is capable of ”generating 4d spacetime” with matter fields, via n isometries which serve as KK-fibres [6]. In addition, the horizon ”generates” axial symmetry as usual.

In more realistic KK theories the n fibre-isometries are replaced by homothetic conformal isometries or still more generally, by periodicity conditions on all fields involved (cf e.g. [1]). This raises the question whether such structures could arise ”automatically” as well, namely from a non-stationary but suitably structured horizon, as a consequence of a hypothetical generalized rigidity theorem. While a concrete implementation of this speculation would lead beyond the scope of the present work, the basic idea does give a motivation for studying general gravitational collapse in higher dimensions, as initiated below.

The key concepts in the study of gravitational collapse are (outer) trapped surfaces and marginally outer trapped surfaces (MOTS). The latter are defined as compact codimension 2-surfaces on which the orthogonally outgoing family of null geodesics has vanishing expansion (cf e.g. [11]). In applications, "stable" and "strictly stable" MOTS play a distinguished role and have been studied extensively [11]–[21]. Here stability is a mild restriction which roughly speaking requires the existence of a codimension 1-neighbourhood in a selected normal direction, which can be foliated by trapped surfaces inside and untrapped surfaces outside. On the other hand, it has been shown that the (global) boundary of the trapped region in a hypersurface is formed by a smooth, strictly stable MOTS [14, 15, 16]. Originally, the stability condition came up in connection with the topology theorems for MOTS [17, 18, 19]. More recently, however, the crucial role of stability in the time evolution of MOTS in a foliated spacetime was clarified, as stability implies smooth evolution while instability signals "jumps" [20]. Moreover, stability is significant for the singularity theorems, as the original requirement of the existence of a trapped surface (both null expansions converging) can in essence be replaced by requirement of existence of a stable MOTS [21]. All these results apply in principle in higher dimensions, (at least till 7), and motivate our interest in the stability condition in particular.

A further motivation comes from the area inequalities for stable MOTS [22, 23]. Such inequalities have been found in particular for axially symmetric, stable 2d MOTS in 4d Einstein-Maxwell by Gabach-Clement, Jaramillo and Reiris [24, 25] and in 4d Einstein-Maxwell dilaton (EMD) theory by Yazadjiev [26]. Other inequalities have been obtained for stable 3d MOTS with various topologies and corresponding symmetries in 5d spactimes by Hollands [27]. Since a 5d vacuum with a KK symmetry is equivalent to EMD theory in 4d, the results of these papers can be compared provided the concepts of stable 2d MOTS and 3d MOTS can be matched. This is the main goal of the present paper.

As we want to illustrate the basic concepts rather than investigating a physically realistic theory, we restrict ourselves to 5d vacuum spacetimes and their dimensional reduction along an isometry. We first (in Sect. 2) review Geroch's projection formalism [28] and continue in Sect. 3.1 with recalling the well-known behaviour of Killing horizons under such a reduction [9, 6]. In Sect. 3.2 we show that 3d MOTS do project down to 2d ones provided an appropriate conformal scaling (which may be ascribed to Pauli [10]) is used for the spacetime metric. The scaling is irrelevant only in a degenerate case in which the norm of the KK Killing field is constant along the outgoing null direction. Fortunately, the Pauli-scaling is the same which is compulsory for matching the variational formulations in different dimensions anyway [1, 29]. Next (in Sect. 3.3) we relate the stability definitions for 3d and 2d MOTS, which is straightforward in principle but involves some subtleties. This allows us (in Sect. 4) to discuss the area inequalities as well. Here we restrict ourselves to relate the limiting cases of the inequalities found by Hollands and Yazadjiev mentioned above.

## 2 Kaluza-Klein a la Geroch

We consider a 5-dimensional Ricci-flat manifold $({}^5\mathcal{M}, {}^5g_{AB})$ of signature (- + + + +), where capital latin indices run from 0 to 4. We assume that there is a spacelike isometry $\mathcal{C}$ without fixed points and a corresponding Killing field $c^A$ on $\mathcal{M}$, i.e. the Lie derivative satisfies $\mathcal{L}_c\ {}^5g_{AB} = 0$. The squared norm of $c^A$ is denoted by $V^2 = c^A c_A > 0$. We assume that the set of orbits of $\mathcal{C}$ is a smooth manifold $\mathcal{M}$. Quantities on ${}^5\mathcal{M}$ are distinguished from the ones on $\mathcal{M}$ either by the superscript ${}^5$, or by capitial indices, or both. A set $\mathcal{E} \subset {}^5\mathcal{M}$ is called **invariant** if $\mathcal{C}(\mathcal{E}) = \mathcal{E}$.

We now recall Geroch's projection formalism [28]; the terms "lift" and "projection" associated with the isometry introduced below are inspired by KK fibre-bundle language [2]; (for sets our terminology is not standard).

**The lift $\mathcal{C}^\uparrow$ of a set $\mathcal{D}$:**
$\mathcal{C}^\uparrow : \mathcal{M} \to {}^5\mathcal{M}$ assigns to a subset $\mathcal{D} \subset \mathcal{M}$ the set of points $\mathcal{E} \subset {}^5\mathcal{M}$ on the orbits through $\mathcal{D}$.

**The projection $\mathcal{C}^\downarrow$ of an invariant set $\mathcal{E}$:**
$\mathcal{C}^\downarrow : {}^5\mathcal{M} \to \mathcal{M}$ assigns to an invariant subset $\mathcal{E} \subset {}^5\mathcal{M}$ the set of orbits $\mathcal{E}/\mathcal{C}$.

These maps induce maps between **invariant tensor fields** $w^{A\ldots M}{}_{N\ldots Z}$ on ${}^5\mathcal{M}$, defined by

$$\mathcal{L}_c w^{A\ldots M}{}_{N\ldots Z} = 0 \quad w^{A\ldots M}{}_{N\ldots Z} c_A = 0 \ \ \ldots \ \ w^{A\ldots M}{}_{N\ldots Z} c^Z = 0 \tag{1}$$

and tensor fields on $\mathcal{M}$. The following maps are of course nothing but "pullbacks" and "push-forwards".


${}^5\mathcal{M}$
$y = \mathcal{C}^\uparrow(x)$
$\mathbb{R}$
$F = \mathcal{C}^\uparrow_*(f)$
$\mathcal{M}$
$x$
$F(y) = f \circ \mathcal{C}^\downarrow(y)$
$f$

**Lift $\mathcal{C}^\uparrow_*$ of a function** $f : \mathcal{M} \to \mathbb{R}; f \in \mathcal{F}(\mathcal{M})$ (the module of functions) **to an invariant function** $F : {}^5\mathcal{M} \to \mathbb{R}; F \in \mathcal{F}({}^5\mathcal{M})$:
$\mathcal{C}^\uparrow_* : \mathcal{F}(\mathcal{M}) \to \mathcal{F}({}^5\mathcal{M})$ is defined as $F = \mathcal{C}^\uparrow_*(f) = f \circ \mathcal{C}^\downarrow$

**Projection of an invariant function** $F \in \mathcal{F}({}^5\mathcal{M})$**:**
$\mathcal{C}^\downarrow_* : \mathcal{F}({}^5\mathcal{M}) \to \mathcal{F}(\mathcal{M})$ is defined as $f = \mathcal{C}^\downarrow_*(F) = F \circ \mathcal{C}^\uparrow$

We will also use this notation for null expansions (cf Sect. 3.2.) which depend on the geometry of submanifolds and are thus not proper functions on $\mathcal{M}$ and ${}^5\mathcal{M}$. The maps introduced above can now be extended straightforwardly to vector and tensor fields by considering gradients of functions etc.; for details cf [28]:

**Lift $\mathcal{C}^{\uparrow}_{*}$ of a vector field $w^i \in T(\mathcal{M})$ to an invariant vector field $z^A \in T({}^5\mathcal{M})$:**
$\mathcal{C}^{\uparrow}_{*} : T(\mathcal{M}) \to T({}^5\mathcal{M})$ denoted by $\mathcal{C}^{\uparrow}_{*}(w^i) = z^A$

**Projection of an invariant vector field $z^A$:**
$\mathcal{C}^{\downarrow}_{*} : T({}^5\mathcal{M}) \to T(\mathcal{M})$ denoted by $\mathcal{C}^{\downarrow}_{*}(z^A) = w^i$

From the construction it is clear that $\mathcal{C}^{\uparrow}$ and $\mathcal{C}^{\downarrow}$ are not isomorphisms, while $\mathcal{C}^{\uparrow}_{*}$ and $\mathcal{C}^{\downarrow}_{*}$ are [28], which is indicated by the asterisks. In Lemma 1 below, the action of these mappings on vector fields will be displayed explicitly in the coordinate system (5) adapted to the isometry.

The projection maps introduced above should not be confused with a particular tensor field called "the projector" defined by

$$P^A_{\ C} = \delta^A_{\ C} - V^{-2} c^A c_C. \tag{2}$$

which can in fact be applied to *any* tensor field in $\otimes T({}^5\mathcal{M})$. In particular, applied to ${}^5g_{AB}$ it gives a metric

$$\widetilde{g}_{CD} = P^A_{\ C} P^B_{\ D}\ {}^5g_{AB}. \tag{3}$$

The covariant derivative w.r.t. ${}^5g_{AB}$ is denoted by ${}^5\nabla$. For an invariant vector field $w^A$ another covariant derivative $\widetilde{\nabla}_A$ can be defined by [28]

$$\widetilde{\nabla}_C w_D = P^A_{\ C} P^B_{\ D}\ {}^5\nabla_A w_B. \tag{4}$$

This derivative $\widetilde{\nabla}_A$ is metric with respect to (3), torsion free and $\widetilde{\nabla}_C w_D$ is still invariant in the sense of (1). Hence it is identical with the covariant derivative on $\mathcal{M}$ with respect to $\widetilde{g}_{CD}$.

Below we will almost exclusively use the conformally rescaled metric $g_{AB} = V\widetilde{g}_{AB}$ on $\mathcal{M}$; in fact this extraction of $V$ will be crucial. The covariant derivative w.r.t $g_{AB}$ is denoted by $\nabla_A$.

A parametrisation of ${}^5g_{AB}$ which will be useful in what follows is

$$ds^2 = {}^5g_{AB} dx^A dx^B = V^2 \left(dx^4 + 2A_i dx^i\right)^2 + V^{-1} g_{ij} dx^i dx^j \tag{5}$$

where latin indices run from 0 to 3 and $V$, $A_i$ and $g_{ij}$ depend on $x^i$ only. Such coordinates exist and will be used below on neighbourhoods $\mathcal{N}$ of compact 2-surfaces $\mathcal{S} \subset \mathcal{M}$ and on the corresponding lift ${}^5\mathcal{N} \supset {}^5\mathcal{S}$. The coordinate $x^4$ is periodic with period $Z$.

Instead of $V$ a more common terminology is either $V = e^{2\varphi/\sqrt{3}}$ [5] or $V = e^{-2\varphi/\sqrt{3}}$ [4], while our definition of $A_i$ follows practice in relativity [4, 5, 6] but differs from practice in gauge theory at least by a factor of 2 [2].

The Lagrangian density on $({}^5\mathcal{M}, {}^5g_{AB})$ can be decomposed as

$$\pounds = \frac{\sqrt{{}^5g}}{16\pi Z}{}^5R = \frac{\sqrt{g}}{16\pi}\left[R - \frac{3}{2}v^{-2}\nabla_i v\nabla^i v - v^3 F_{ij}F^{ij}\right] \tag{6}$$

where $v = \mathcal{C}_*^{\downarrow}(V)$ and $F_{ij} = 2\nabla_{[i}A_{j]}$. This Lagrangian determines EMD theory on $(\mathcal{M}, g_{ij})$. The metric variable $A_i$ in (5), which is the dynamic variable in (6) can be gauge transformed via $A'_i = A_i + \nabla_i\Lambda$ for some function $\Lambda$.

We now give the action of $\mathcal{C}_*^{\uparrow}$ explicitly in the coordinates (5).

**Lemma 1**. Let $w^i$ be a vector field on $(\mathcal{M}, g_{ij})$. On $({}^5\mathcal{M}, g_{AB})$ and in adapted coordinates (5), we define $w^A = (w^i, 0)$ and $\alpha = 2w^iA_i$. Then the lift $\mathcal{C}_*^{\uparrow}(w^i)$ is given in these coordinates by

$$\mathcal{C}_*^{\uparrow}(w^i) = z^A := w^A - \mathcal{C}_*^{\uparrow}(\alpha)c^A \tag{7}$$

**Proof.** It is easy to see that the vector field $z^A$ is invariant, viz. $z^Ac_A = 0$ and $\mathcal{L}_c z^A = 0$, and it obviously projects down to $w^i$, i.e. $\mathcal{C}_*^{\downarrow}(z^A) = w^i$. The assertion now follows since $\mathcal{C}_*^{\uparrow}$ is an isomorphism.

**Remark.** The preceding presentation has a counterpart in the fibre bundle formulation of gauge theory [2]. In the general setting, the fibres are not necessarily isometries. This entails, in particular, that there is *a priori* no natural "lift" of a vector field, but any connection on the bundle specifies a "horizontal lift". In the present case a natural conncection is provided by Killing transport, and our lift is indeed "horizontal" in this sense. What we have done in (7) now reads, in gauge theoretic language, to decompose the horizontal lift of a vector field in terms of a "direct product basis" (5). Applied to the basis vectors themselves, this corresponds to the introduction of a "gauge covariant derivative" ( cf Sect. 5 of [2]).

# 3 Horizons

## 3.1 Killing horizons

Let $l^i$ be a Killing field on $\mathcal{M}$ and $\mathcal{H} \subset \mathcal{M}$ a Killing horizon. This means that $\mathcal{H}$ is a 3d null surface with "generator" $l^i$ which is null on $\mathcal{H}$ and satisfies $l^i\nabla_i l^j = \kappa l^j$ there, where $\kappa$ is the surface gravity which is constant on the horizon (for this and the following statements, cf Carter [30]). As $l^i$ is only fixed up to a constant, the same is true for $\kappa$, but conditions at spatial infinity can remove this freedom. In the presence of a Maxwell (and possibly dilaton) field, so in particular for the EMD Lagrangian (6), $v$ and $\varphi = 2A_il^i$ are constant on the horizon as well. The latter constant is again arbitrary, as there is still the rescaling freedom in $l^i$ and in addition the gauge freedom $A'_i = A_i + \nabla_i\Lambda$. However, as above asymptotic conditions (namely falloff for $A_i$) can fix $\varphi$ uniquely.

We now recall (and reformulate) the following result (cf Sect. 4 of [6]):

**Theorem 1**. In an Einstein-Maxwell-dilaton spacetime $(\mathcal{M}, g_{ij})$ let $\mathcal{H}$ be a Killing horizon with generator $l^i$ and surface gravity $\kappa$, i.e. $l^i\nabla_i l^j = \kappa l^j$ on $\mathcal{H}$. Then the lift ${}^5\mathcal{H} = \mathcal{C}^\uparrow(\mathcal{H})$ is a Killing horizon in the 5d vacuum spacetime $({}^5\mathcal{M}, {}^5g_{AB})$ with generator $\mathcal{C}^\uparrow_*(l^i) = h^A = l^A - \Phi c^A$ where $\varphi = 2A_i l^i$ and $\Phi = \mathcal{C}^\uparrow_*(\varphi)$. Moreover, the surface gravity $\kappa$ is as before, i.e. $h^A\,{}^5\nabla_A h^B = \mathcal{K}h^B$ on ${}^5\mathcal{H}$ with $\mathcal{K} = \mathcal{C}^\uparrow_*(\kappa)$

**Proof**. We first note that $h^A$ is null on ${}^5\mathcal{H}$, which is obvious either from

$$\mathcal{C}^\downarrow_*\left(h^A h_A\right) = \mathcal{C}^\downarrow_*\left(h^A\right)\mathcal{C}^\downarrow_*\left(h_A\right) = l^i l_i = 0 \tag{8}$$

as all objects are invariant, or in coordinates via

$$\mathcal{C}^\downarrow_*\left(h^A h_A\right) = v^2\left(\varphi - 2A_i l^i\right)^2 + v^{-1}g_{ij}l^i l^j = 0. \tag{9}$$

The non-trivial statements of the theorem now follow from

$$h^A\,{}^5\nabla_A h_B = -\frac{1}{2}\,{}^5\nabla_B(h^A h_A) = -\frac{1}{2}\mathcal{C}^\uparrow_*\left(\nabla_i(v^{-1}l^j l_j)\right) = \mathcal{C}^\uparrow_*\left(v^{-1}l^j\nabla_j l_i\right) = \mathcal{K}\mathcal{C}^\uparrow_*\left(v^{-1}l_i\right) = \mathcal{K}h_B \tag{10}$$

on ${}^5\mathcal{H} = \mathcal{C}^\uparrow(\mathcal{H})$, where we have used adapted coordinates (5) and Eqn. (9).

**Remark.** The notation above, in particular regarding the constants $\Phi = \mathcal{C}^\uparrow_*(\varphi)$ and $\mathcal{K} = \mathcal{C}^\uparrow_*(\kappa)$ may appear pedantic. It is adopted to be consistent with calculations below where analogous functions on ${}^5\mathcal{M}$ are used which are either no longer constant, or (in Theorem 3) not even invariant. There, care with notation pays off to recognize subtleties.

## 3.2 The expansion

We now consider a spacelike 2-surface $\mathcal{S} \subset \mathcal{M}$, and null vectors $l^i$ and $k^i$, not necessarily Killing but orthogonal to $\mathcal{S}$ and scaled such that $l^i\ k_i = -2$. These vectors can be extended off $\mathcal{S}$ by considering the null geodesics with these tangents. This allows to define the expansion of $l^i$ (or of the corresponding null geodesics) on $\mathcal{S}$ via

$$\Theta = \left(g^{ij} + l^{(i}k^{j)}\right)\nabla_i l_j. \tag{11}$$

A standard result (cf e.g. Theorem 3.6.1 of [31]) reads that $\Theta$ is in fact independent of the way how $l^i$ has been extended. For EMD theory (6) we define $\psi = 2A_i k^i$ and we recall that $\varphi = 2A_i l^i$. These quantities will in general not be constant on $\mathcal{S}$. We now lift $\mathcal{S}$ to a 3-surface ${}^5\mathcal{S} = \mathcal{C}^\uparrow(\mathcal{S})$, define $\Phi = \mathcal{C}^\uparrow_*(\varphi)$ and $\Psi = \mathcal{C}^\uparrow_*(\psi)$, and lift $l^i$ and $k^i$ to invariant null vectors $\mathcal{C}^\uparrow_*(l^i) = h^A = l^A - \Phi c^A$ and $\mathcal{C}^\uparrow_*(k^i) = j^A = k^A - \Psi c^A$ on ${}^5\mathcal{M}$. It follows that $g_{AB}h^A j^B = -2V^{-1}$, and in analogy with (11) we define the expansion of $h^A$:

$${}^5\Theta = \left(g^{AB} + Vh^{(A}j^{B)}\right){}^5\nabla_A h_B. \tag{12}$$


$l^A = (l^i, 0)$
${}^5\mathcal{S} = \mathcal{C}^{\uparrow}(\mathcal{S})$
$h^A = \mathcal{C}_*^{\uparrow}(l^i)$
$c^A$
$k^A = (k^i, 0)$
$j^A = \mathcal{C}_*^{\uparrow}(k^i)$
${}^5\mathcal{M}$
$l^i$
$\mathcal{S}$
$\mathcal{M}$
$k^i$

Alternatively, we could of course start from a 3d surface in ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ which is invariant under an isometry. It projects to a 2d surface $\mathcal{S} \subset \mathcal{M}$, and the null normals and expansions are related as above.

We now have the following result

**Theorem 2**. Let $\mathcal{S} \subset \mathcal{M}$ and ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ be spacelike 2-surfaces such that $\mathcal{S} = \mathcal{C}^{\downarrow}({}^5\mathcal{S}) = {}^5\mathcal{S}/\mathcal{C}$ where $\mathcal{C}$ is an isometry. Let $l^i$ and $h^A = \mathcal{C}_*^{\uparrow}(l^i)$ be the respective null normals related by the Geroch-isomorphism. Then the corresponding null expansions defined by (11) and (12) coincide in the sense that $\mathcal{C}_*^{\downarrow}\left({}^5\Theta(h)\right) = \Theta(l)$.

**Proof.** For ease of notation, we suppress the $\mathcal{C}_*^{\downarrow}-$ notation in the subsequent calculation. However, we have to take care to avoid ill-defined expressions such as $\mathcal{C}_*^{\downarrow}\left(g^{AB} + Vh^{(A}j^{B)}\right)$ and $\mathcal{C}_*^{\downarrow}\left({}^5\nabla_A h_B\right)$, which is accomplished by appropriate algebraic decomposition of ${}^5\nabla_A h_B$. Next, to obtain (16) we have used (4) while the final step which leads to (19) uses the behaviour of the 4d covariant derivative under conformal rescalings of the metric.

$$
\begin{aligned}
{}^5\Theta(h) &= \left(g^{AB} + Vh^{(A}j^{B)}\right) {}^5\nabla_A h_B = &(13)\\
&= \left(g^{AB} + Vh^{(A}j^{B)}\right)\left(P_A^C + V^{-2}c_A c^C\right)\left(P_B^D + V^{-2}c_B c^D\right) {}^5\nabla_C h_D = &(14)\\
&= \left(g^{AB} + Vh^{(A}j^{B)}\right) P_A^C P_B^D\, {}^5\nabla_C h_D + V^{-2}c^A c^B\, {}^5\nabla_A h_B = &(15)\\
&= \left(\widetilde{g}^{ij} + vl^{(i}k^{j)}\right)\widetilde{\nabla}_i(V^{-1}l_j) - v^{-2}c^A h^B\, {}^5\nabla_A c_B = &(16)\\
&= v\left(g^{ij} + l^{(i}k^{j)}\right)\widetilde{\nabla}_i(v^{-1}l_j) + v^{-2}c^A h^B\, {}^5\nabla_B c_A = &(17)\\
&= \left(g^{ij} + l^{(i}k^{j)}\right)\widetilde{\nabla}_i l_j + v^{-1}l^i\nabla_i v = &(18)\\
&= \left(g^{ij} + l^{(i}k^{j)}\right)\nabla_i l_j - v^{-1}l^i\nabla_i v + v^{-1}l^i\nabla_i v = \Theta(l) &(19)
\end{aligned}
$$

**Remark.** The correct conformal scaling as in (5) is crucial both for the Kaluza-Klein decomposition (6) as well as for the previous result. Only in the special case where $l^i$ leaves $v$ invariant, i.e. $l^i\nabla_i v = 0$, the conformal scaling is irrelevant.

### 3.3 MOTS

A **MOTS** $\mathcal{S} \subset \mathcal{M}$ is a compact, embedded 2-surface with vanishing null expansion, i.e. $\Theta(l) = 0$. Analogously, A **MOTS** ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ is a compact embedded 3-surface with ${}^5\Theta(h) = 0$.

Theorem 2 shows that a MOTS $\mathcal{S}$ can be lifted naturally, while an invariant MOTS ${}^5\mathcal{S}$ can be projected. In view of the significance of the property of stability as outlined in the introduction, we wish to show that it is preserved upon projections and lifts as well. In [12, 13] two definitions for (strict) stability of MOTS were given. We recall below, for 2d MOTS $\mathcal{S} \subset \mathcal{M}$, the definition which was called "(strictly) stably outermost" in [12, 13] and discuss its extension to 3d MOTS ${}^5\mathcal{S} \subset {}^5\mathcal{M}$. The alternative definition required non-negativity (positivity) of the principal eigenvalue of a linear elliptic "stability operator". These definitions are equivalent, and this equivalence will play a role in Theorem 3 below.

**Stability in 4d:** A MOTS $\mathcal{S} \subset \mathcal{M}$ is called stable w.r.t. a normal direction $m^i$ iff there exists a variation $\delta_{\psi m}$ with $\psi \not\equiv 0$ such that $\delta_{\psi m}\Theta \geq 0$. The MOTS is strictly stable iff it is stable with $\delta_{\psi m}\Theta \not\equiv 0$.

**Stability in 5d:** A MOTS ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ is called stable w.r.t. a normal direction $n^A$ iff there exists a variation $\delta_{\Psi n}$ with $\Psi \not\equiv 0$ such that $\delta_{\Psi n}{}^5\Theta \geq 0$. The MOTS is strictly stable iff it is stable with $\delta_{\Psi n}{}^5\Theta \not\equiv 0$.

In principle any normal directions $m^i$ or $n^A$ can be chosen here. For $m^i = l^i$ or $n^A = h^A$ (the null vectors defining the MOTS) the variations considered above reduce to the Raychaudhuri equation and the stability conditions become highly restrictive. The least restrictive choices are $m^i = k^i$ or $n^A = j^A$ (the null directions *not* defining the MOTS).

We now relate stabilty for 2d and 3d MOTS. *A priori* a MOTS ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ need not be invariant, nor need the direction of stability be invariant under the isometry. On the other hand, there *is* a relation between stability and the action of an isometry: If the MOTS is stable in a normal direction $n^A$, then any Killing field $d^A$ is either tangent to the MOTS, or its normal component $d^A_\perp$ necessarily lies in the conical segement between the null generator $h^A$ of the MOTS and $n^A$ (cf. Theorem 8.1. of [13] and the subsequent discussion). However, to keep the theorem below simple, we assume in its second part invariance of both the MOTS itself as well as of the stability direction.

**Theorem 3.**

1. Let $\mathcal{S} \subset \mathcal{M}$ be a MOTS which is (strictly) stable w.r.t. a normal direction $m^i$. Then the lifted MOTS $\mathcal{C}^\uparrow(\mathcal{S}) \subset {}^5\mathcal{M}$ is (strictly) stable w.r.t. the normal direction $n^A = \mathcal{C}^\uparrow_*(m^i)$.

2. Let ${}^5\mathcal{S} \subset {}^5\mathcal{M}$ be an invariant MOTS which is (strictly) stable w.r.t. the invariant normal direction $n^A$. Then the projected MOTS $\mathcal{S} = \mathcal{C}^{\downarrow}({}^5\mathcal{S}) \subset \mathcal{M}$ is (strictly) stable w.r.t. the normal direction $m^i = \mathcal{C}^{\downarrow}_{*}(n^A)$.

**Proof**.

1. This part is trivial: Let $\psi$ be the function in the definition of stability of $\mathcal{S} \subset \mathcal{M}$ such that $\delta_{\psi m}\Theta \geq 0$. Then the lifts $\Psi = \mathcal{C}^{\uparrow}_{*}\psi$ and $n^A = \mathcal{C}^{\uparrow}(m^i)$ satisfy the requirements for stability on ${}^5\mathcal{S} \subset {}^5\mathcal{M}$, in particular $\delta_{\Psi n}{}^5\Theta \geq 0$.

2. In spite of the invariance assumptions there is still a subtlety here: The function $\Psi$ which defines the variation $\delta_{\Psi n}{}^5\Theta \geq 0$ in the definition of stability of ${}^5\mathcal{S}$ need not be invariant under $\mathcal{C}$, in which case it does not project to $\mathcal{S} \subset \mathcal{M}$. Here we need the elliptic machinery developed in [12, 13]. We rewrite the variation $\delta_{\Psi n}{}^5\Theta$ in terms of a linear elliptic operator ${}^5L_n$ acting on $\Psi$ such that $\delta_{\Psi n}{}^5\Theta = {}^5L_n\Psi$. Any linear elliptic operator has a real, positive "principal" eigenfunction $\Phi$ corresponding to the "principal" eigenvalue $\lambda$ (the eigenvalue with lowest real part), viz. ${}^5L_n\Phi = \lambda\Phi$. For ${}^5L_n$ introduced above, the definitions of stability and strict stability are equivalent to $\lambda \geq 0$ and $\lambda > 0$, respectively. This means in particular that for a stable MOTS, the principal eigenfunction $\Phi$ defines a prefered class of variations such that $\delta_{\Phi n}{}^5\Theta = {}^5L_n\Phi = \lambda\Phi \geq 0$. (or $> 0$ in the strictly stable case). But by virtue of the invariance of the stability direction $n^A$, the stability operator commutes with the isometry, viz. $\mathcal{L}_c L_n = L_n \mathcal{L}_c$. Now Theorem 8.2. of [13] implies that the principal eigenfunction is invariant under the isometry, i.e. $\mathcal{L}_c\Phi = 0$. Hence $\Phi$ can be projected to $\mathcal{M}$ and $\phi = \mathcal{C}^{\downarrow}_{*}\Phi$ defines the variations on $\mathcal{S}$ which imply stability, in particular $\delta_{\phi n}\Theta \geq 0$.

# 4 Area inequalities

Area inequalities bound the area $A$ of stable MOTS in terms of quantities defined on the MOTS, such as an electric charge $Q$, a magnetic charge $P$ and, in the axially symmetric case, the angular momentum $J$ ([22, 23]). Here we briefly recall and compare the area inequalities for Einstein-Maxwell, Einstein-Maxwell-dilaton and 5d vacuum.

For axially symmetric, stable 2d MOTS in Einstein-Maxwell theory, Gabach-Clement et al. [24, 25] proved that

$$A \geq 4\pi\sqrt{4J^2 + (Q^2 + P^2)^2} \tag{20}$$

where equality holds iff the near horizon geometry is an extreme Kerr-Newman one.

In EMD theory, for couplings which include Einstein-Maxwell (i.e. no dilaton) and the Lagrangian (6), Yazadjiev [26] has shown that for stable 2d MOTS

$$A \geq 8\pi\sqrt{|J^2 - Q^2P^2|} \tag{21}$$

Moreover, equality holds iff the coupling is the KK one (6) and the near horizon geometry is extreme and stationary.

On the other hand, in a 5-dimensional vacuum spacetime with isometry group $U(1) \times U(1)$ Hollands [27] has obtained an area inequality of stable 3d MOTS ${}^5\mathcal{S}$ in the form

$$
{}^5A \geq 8\pi\sqrt{|{}^5J_1\ {}^5J_2|} \tag{22}
$$

where ${}^5J_1$ and ${}^5J_2$ are angular momenta associated with the isometries. Again equality holds precisely for the stationary, extreme (near-) horizon geometries which have been studied in detail [33, 34].

We now compare these inequalities. Regarding the relation between (20) and (21) the situation is clear: As already noticed in [26], the former is the stronger inequality but it only applies in the Einstein-Maxwell case. On the other hand, (21) and (22) should have a common range of applicability: A 5d vacuum with a spacelike isometry is equivalent to EMD theory in 4d, and the isometry preserves stable MOTS in the sense of Theorems 2 and 3.

Rather than proving equivalence of the inequalities (21) and (22) we restrict ourselves here to examining the respective limiting cases. As mentioned above, (22) is saturated iff the "near-horizon-geometry" is extreme and stationary, and such geometries have been investigated thoroughly [33, 34]. It has been shown that the possible horizon topologies take one of the forms $S^3$, $S^2 \times T$, $T^3$, or the lens spaces $L(p, q)$, and $T^3$ can be excluded under mild extra assumptions [35]. On the other hand, (21) is only valid for MOTS of spherical topology. Thus a prerequisite for matching (21) with (22) is that the MOTS ${}^5\mathcal{S}$ is a bundle over $S^2$. This restricts the possible candidates to $S^3$, $S^2 \times T$ and $S^3/\mathbb{Z}_p = L(p, 1)$ [36] but excludes the Lens spaces $L(p, q)$ for $q \geq 2$. However, as the latter *are* covered by (22), this inequality is *a priori* more general than (21).

In the cases of $S^2$ bundles, all extreme stationary horizon geometries arise from the explicitly known 4-parameter family of stationary KK black holes [7, 8], either by restricting parameters or by performing appropriate limits (cf Appendix A1 of [32]). For dimensional reduction, there arises in each case an isometry without fixed points as natural candidate for a fibre. The corresponding bundle can be classified by the degree $p$ of the fibration, which for the known extreme solutions is related to the constant length $Z$ of the fibres and the magnetic monopole via $8\pi P = pZ$. In the case of $S^2 \times T$ topology the bundle is trivial which means $P = 0$. The resulting geometries include the boosted extremal Kerr and the black ring ones, and the identification of (21) and (22) is accomplished via $J_1 = J_2 = ZJ$ and ${}^5A = ZA$.

In the case of $S^3$ the geometry of the MOTS can be identified either as extreme Myers-Perry or as extremal rotating Kaluza-Klein solution. In these cases, the canonical Killing vectors of Thm. 1 of [27] cannot be used for reduction as they have zeros. Rather, one has to take appropriate linear combinations, and the corresponding ("Komar"-) momenta lead to the identification of (21) with (22) via $J_1 = Z(J - PQ)$ and $J_2 = Z(J + PQ)$ (for details cf Appendix A.2. of [32]).

We finally remark that, from Theorem 1, the property of extremality ($\kappa = 0$) of a stationary Killing horizon is preserved upon lift and projection, irrespective of any additional symmetries. This behaviour is of course reflected in the axially symmetric cases discussed above.

**Acknowledgements.** We are grateful to Piotr Bizoń, Piotr Chruściel, Marc Mars and Helmuth Urbantke for helpful discussions.

# References

[1] Overduin J M and Wesson P S 1997 *Phys. Rep.* **283** 303

[2] Cho Y M 1975 *J. Math. Phys.* **16** 2029

[3] Emparan R and Reall H S 2008 *Living Rev. Rel.* **11:6**

[4] Horowitz G T and Wiseman T *in* 2012 Horowitz G T (ed.) *Black Holes in Higher Dimensions* (Cambridge: Cambridge University Press)

[5] Gibbons G 1982 *Nucl. Phys. B* **207** 337

[6] Simon W 1985 *Gen. Rel. Grav.* **17** 439

[7] Rasheed D 1995 *Nucl. Phys. B* **454** 379

[8] Larsen F 2000 *Nucl. Phys. B* **575** 211

[9] Hollands S and Ishibashi A 2012 *Class. Quantum Grav.* **29** 163001

[10] Pauli W *in* 1955 Jordan P *Schwerkraft und Weltall* (Braunschweig: F. Vieweg und Sohn)

[11] Jaramillo J L 2011 *Int. J. of Modern Physics* **20** 2169

[12] Andersson L, Mars M and Simon W 2005 *Phys. Rev. Lett.* **95** 111102

[13] Andersson L, Mars M and Simon W 2008 *Adv. Theor. Math. Phys* **12** 853

[14] Andersson L and Metzger J 2009 *Commun. Math. Phys* **290** 941

[15] Eichmair M 2009 *J. Diff. Geom* **83** 551

[16] Andersson L, Eichmair M and Metzger J *in* 2009 Agranovsky M et al. (eds) *Proceedings of the Fourth International Conference on Complex Analysis and Dynamical Systems (Contemporary Mathematics 553)* (Providence: American Mathematical Society and Ramat-Gan: Bar-Ilan University)

[17] Hawking S W and Ellis G F R 1973 *The large scale structure of space-time* (Cambridge: Cambridge University Press).

[18] Newman R P A C 1987 *Class. Quantum Grav.* **4** 277

[19] Galloway G *in* 2012 Horowitz G T (ed.) *Black Holes in Higher Dimensions* (Cambridge: Cambridge University Press)

[20] Andersson L, Mars M, Metzger J and Simon W 2009 *Class. Quantum Grav.* **26** 085018

[21] Costa e Silva I P 2012 *Class. Quantum Grav.* **29** 235008

[22] Dain S 2012 *Class. Quantum Grav.* **29** 073001

[23] Jaramillo J L 2013 *Springer Proceedings in Mathematics & Statistics* **26** 139

[24] Gabach-Clement M E and Jaramillo J L 2012 *Phys. Rev. D 86* 064021

[25] Gabach-Clement M E, Jaramillo J L and Reiris M 2013 *Class. Quantum Grav.* (to be published)

[26] Yazadjiev S 2013 *Phys. Rev. D* **87** 024016

[27] Hollands S 2012 *Class. Quantum Grav.* **29** 065006

[28] Geroch R 1971 *J. Math. Phys.* **12**, 918

[29] Cho Y M 1992 *Phys. Rev. Lett.* **68** 3133

[30] Carter B *in* 1973 DeWitt C & DeWitt B S (eds.) *Black hole equilibrium states* (New York: Gordon & Breach)

[31] Jost J 1998 *Riemanian Geometry and Geometric Analysis* (Berlin, Heidelberg: Springer)

[32] Emparan R and Maccarrone A 2007 *Phys Rev. D* **75** 084006

[33] Kunduri H K and Lucietti J 2009 *J. Math. Phys.* **50** 082502

[34] Hollands S and Ishibashi A 2010 *Ann. Henri Poincaré* **10** 1537

[35] Holland J 2010 http://arxiv.org/abs/1008.0520 (unpublished)

[36] Steenrod N 1944 *The Annals of Mathematics, 2nd Ser.* **45** 294